for the ages 20 to 50 compare favourably with the higher among New Zealand societies. This last made from the limited data, it is most probable that, ill show a very considerable comparative increase.	Average Number of Weeks' Sickness per Annum to each Member actually sick.
of weeks' sickness <i>per member</i> during the decade f iss experienced <i>per member sick</i> is considerably as been said, it is not safe to rely on deductions $n$ d, the sickness experience at the higher ages wi	Average Number of Weeks' Sickness per Annum to each Member.
The percentage of members sick and also the average number of weeks' sickness <i>per member</i> during the decade for the ages 20 to 50 compare favourably with the English experience, whereas the average number of weeks' sickness experienced <i>per member sick</i> is considerably higher among New Zealand societies. This last observation applies especially to the higher ages; and, although, as has been said, it is not safe to rely on deductions made from the limited data, it is most probable that, if there is in the colony a greater average longevity than in England, the sickness experience at the higher ages will show a very considerable comparative increase. The respective results are as follow :	Percentage of Members sick during a Year.

	Perc	Percentage of Members sick during a Year.	sick during a I	ſear.	Average.	Average Number of Weeks' Sickness per Annum to each Member.	Sickness per Aı nber.	num to	Average Nu	Average Number of Weeks' Sickness per Annum to each Member actually sick.	kness per Ann ally sick.	am to each
Ages of Members.		New Zealand.		Encland.	-	New Zealand.		England.		New Zealand.		England,
	Quinquennium, 1873-77.	Quinquennium, 1878-82.	Decade, 1873-82.	A.O.F., 1871-75.	Quinguennium, 1873–77.	Quinquennium, 1878-82.	Decade, 1873-82.	A.Ö.F., 1871–75.	Quinquennium, 1873-77.	Quinquennium, 1878-82.	Decade, 1873-82.	A.O.F. 1871-75.
Under 20	[1-61]	[15.55]	[18.00]	22-9	[0.87]	[0.42]	[0-83]	1-04	[4.55]	[4.79]	[4.62]	:
<b>2</b> 0 to 25	12.2	13.05	12.69	21.5	0.52	75-0	0.55	0-82	4.29	4.33	4·31	3.57
25 to 30	12.0	11.53	11.72	21.6	0-57	0-63	0.60	0.85	4.74	5.45	5.16	3-96
<b>3</b> 0 to 35	13.8	12.93	13.38	22.3	0-74	0.72	0.73	16-0	5.35	5.58	5.46	4.48
35 to 40	14-7	14.83	14.77	23.4	0-98	1.03	1.01	1.15	6.68	6-97	6-84	Q1.Q
40 to 45	15.3	15.15	15.21	25.0	1.20	1.15	1.17	1.37	7.82	1.61	7.69	<b>5</b> .86
45 to 50	14·1	15.86	15.40	26.9	0.86	1.52	1.33	17-1	6.13	19-6	8-80	6.83
<b>5</b> 0 to 55	[21•4]	[20.17]	[20.36]	29.6	[2.68]	[1.72]	[1.87]	2.26	[12-50]	[8:55]	[9.19]	8-42
55 to 60	[28-8]	[24.62]	[25.34]	34.0	[2.76]	[4.34]	[4-07]	3.21	[9.58]	[17.61]	[16-04]	10-86
60 to 65	[38•4]	[33.53]	[35.29]	43.3	[9-32]	[00.9]	[6-57]	4.59	[24.28]	[14.90]	[18-61]	13.50
65 to 70	[74·1]	[40.51]	[49-06]	52.3	[26.35]	[6:48]	[11.55]	7.97	[35.57]	[16-01]	[23.54]	18-39
70 and upwards	[94.1]	[49.06]	[60.00]	60.1	[39-23]	[20.60]	[25.12]	13.48	[41.68]	[41.99]	[41-87]	24-85
Ages unspecified	[6-1]	[17·86]	[14·29]	•	[0-28]	[2.10]	[1.55]	:	[4·62]	[11-76]	[10-83]	:
All ages	14.1	14.28	14.19	23.45	0-87	1.01	0-95	1.36	6-20	7.10	6.72	5.79

## **H**.—5.