

*The Prevalence of Sickness and Death from Infectious Diseases throughout the Otago-Southland District.—Returns for the Years subsequent to Compulsory Notification.*

Year.	Scarlet Fever or Scarlatina.		Diphtheria.		Enteric or Typhoid Fever.		Tuberculosis.		Puerperal Fever.*		Totals for all Notifiable Infectious Diseases: Cases.
	Number of Cases notified.	Case-rate per 1,000 Population.	Number of Cases notified.	Case-rate per 1,000 Population.	Number of Cases notified.	Case-rate per 1,000 Population.	Number of Cases notified.	Case-rate per 1,000 Population.	Number of Cases notified.	Case-rate per 1,000 Population.	
1901-2 ..	10	0.063	32	0.20	7	0.02	149	0.90	..	..	198
1902-3 ..	843	5.25	36	0.226	24	0.05	160	1.006	13	0.08	1,076
1903-4 ..	510	3.21	12	0.08	42	0.26	65	0.409	3	0.019	632
1904-5 ..	126	0.80	29	0.12	43	0.26	86	0.541	11	0.07	295
1905-6 ..	136	0.85	25	0.16	28	0.18	102	0.642	10	0.063	301
1906-7 ..	180	1.13	23	0.15	23	0.15	95	0.60	10	0.063	331
1907-8 ..	246	1.50	58	0.35	36	0.217	152	0.90	33	0.14	525
1908-9 ..	606	3.65	84	0.50	25	0.15	170	1.02	49	0.30	934
1909-10 ..	367	2.04	95	0.53	71	0.402	134	0.75	14	0.08	681
1910 ..	254	1.42	84	0.47	84	0.47	138	0.778	9	0.05	569

\* Puerperal fevers: The figures prior to 1910 include cases of general blood-poisoning as well as puerperal fevers. Figures for 1910 are for puerperal fevers only.

The following table shows the sexes of cases notified, and the deaths from notifiable infectious diseases in the Otago-Southland District.

Notifiable Diseases.	At all ages.			
	Cases.		Deaths.	
	M.	F.	M.	F.
Scarlet fever .. .. .	117	137	0	0
Diphtheria .. .. .	41	43	3	2
Enteric or typhoid fever .. .. .	57	27	2	4
Tuberculosis .. .. .	61	77	47	60
Puerperal fevers .. .. .	0	9	0	5
Totals .. .. .	276	293	52	71
	569		123	