

TABLE A.—NOTIFIABLE DISEASES IN NEW ZEALAND, 1928, SHOWING DISTRIBUTION BY MONTHS.

Month.	Scarlet Fever.	Diphtheria.	Enteric Fever.	Tuberculosis.	Cerebro-spinal Meningitis.	Poliomyelitis.	Pneumonic Influenza.	Pneumonia.	Mysipelas.	Puerperal Fever.	Septic Abortion.	Relapsing.	Tetanus.	Hydatids.	Trachoma.	Ophthalmia Neonatorum.	Lethargic Encephalitis.	Food Poisoning.	Chronic Lead Poisoning.	Dysentery.	Actinomycosis.	Leprosy.	Totals.
January	201	80	29	85	2	5	23	67	17	7	6	4	..	4	2	1	2	2	1	1	539
February	204	40	19	163	1	5	8	49	22	18	9	3	1	4	6	552
March	428	81	32	133	..	11	11	48	23	12	9	4	1	6	1	2	1	1	..	11	1	..	816
April	529	132	38	92	3	11	10	76	26	12	9	4	1	8	4	..	4	2	..	3	964
May	567	167	48	150	3	4	24	79	26	17	9	7	..	3	1	3	1	6	..	2	1	..	1,118
June	652	236	15	131	1	1	33	104	35	11	4	9	..	5	2	3	4	1	..	1	1	..	1,249
July	798	209	18	114	43	135	46	22	10	9	1	1	..	3	4	..	1	3	1,417
August	754	189	23	171	1	1	66	199	42	19	12	9	4	4	1	3	4	1,500
September	527	136	11	135	5	3	38	138	43	19	10	11	1	9	..	1	2	1	1,090
October	513	103	20	138	2	2	36	156	49	14	8	14	3	11	..	6	2	..	3	1	1,081
November	499	132	20	108	3	2	37	131	33	14	8	4	2	4	1	1	4	6	1	1,010
December	455	95	17	92	2	2	25	103	23	19	19	10	4	3	1	3	873
Totals	6,127	1,600	290	1,512	23	47	354	1,285	385	184	113	88	18	62	13	24	27	19	6	27	4	1	12,209