

TABLE A.—NOTIFIABLE DISEASES IN NEW ZEALAND, 1931, SHOWING DISTRIBUTION BY MONTHS.

Month.	Scarlet Fever.	Diphtheria.	Enteric Fever.	Tuberculosis.	Cerebro-spinal Meningitis.	Polymyelitis.	Influenza.	Dysipelas.	Puerperal Fever.		Eclampsia.	Tetanus.	Hydatids.	Trachoma.	Ophthalmia Neonatorum.	Lethargic Incephalitis.	Food Poisoning.	Bacillary Dysentery.	Leprosy.	Undulant Fever.	Actinomycosis.	Anthrax.	Beriberi.	Chronic Lead Poisoning.	Totals.	
									Ordinary.	Following Abortion or Miscarriage.																
January ..	85	57	21	92	2	2	3	22	11	20	7	6	5	..	7	2	5	2	849
February ..	109	62	13	96	..	1	..	13	13	17	10	4	7	..	1	2	352	
March ..	101	106	17	96	2	2	2	23	22	15	6	2	3	..	2	2	409	
April ..	127	126	8	90	2	1	1	17	9	12	7	1	3	..	4	416	
May ..	127	163	11	76	3	..	1	22	10	2	5	..	3	..	1	439	
June ..	118	119	9	74	2	1	3	33	12	7	5	1	3	..	2	398	
July ..	121	120	17	95	4	4	5	21	16	16	11	1	3	..	1	441	
August ..	125	184	6	81	1	2	4	22	15	6	8	..	5	..	3	462	
September ..	107	108	9	113	1	3	103	14	15	7	5	..	3	..	3	499	
October ..	101	74	11	113	2	4	96	24	16	10	12	..	4	..	1	478	
November ..	76	84	12	94	2	4	16	14	8	14	6	4	8	..	2	..	13	380	
December ..	107	124	27	89	2	1	9	8	13	7	2	2	4	..	5	..	4	410	
Totals ..	1,304	1,327	161	1,109	22	25	247	233	160	133	84	21	59	9	33	12	24	31	1	13	1	1	1	2	5,013	