

TABLE A.—NOTIFIABLE DISEASES IN NEW ZEALAND, 1932, SHOWING DISTRIBUTION BY MONTHS.

Month.	Scarlet Fever.	Diphtheria.	Enteric Fever.	Tuberculosis.	Cerebro-spinal Meningitis.	Poliomyelitis.	Influenza.	Myxipelas.	Puerperal Fever.				Tetanus.	Hydatids.	Trachoma.	Ophthalmia Neonatorum.	Lethargic Encephalitis.	Food Poisoning.	Bacillary Dysentery.	Actinomycosis.	Undulant Fever.	Lead Poisoning.	Leprosy.	Anthrax.	Beriberi.	Phosphorus Poisoning.	Pneumonia.	Totals.
									Ordinary.	Following Abortion or Miscarriage.	Relapsing.																	
January ..	79	62	32	66	..	12	1	7	3	11	8	5	4	..	3	1	6	7	1	..	1	309
February ..	68	52	16	95	..	30	3	21	6	10	3	2	3	..	3	4	4	..	2	322	
March ..	81	88	18	77	3	39	2	20	3	15	4	1	4	1	2	1	4	..	1	375	
April ..	81	74	21	58	1	23	..	23	11	10	8	2	2	..	1	3	12	..	2	333	
May ..	75	78	14	76	2	8	4	22	19	6	7	2	2	..	3	2	1	..	1	1	323	
June ..	72	89	15	79	..	6	5	22	9	8	6	..	3	1	1	1	12	..	1	2	332	
July ..	81	73	19	89	2	1	1	21	9	11	6	2	2	..	1	313	
August ..	68	68	10	52	..	8	3	19	13	17	6	1	2	1	..	3	2	..	2	277	
September ..	55	58	5	81	2	5	2	24	15	19	2	2	5	1	3	279	
October ..	52	51	9	101	2	2	1	17	9	12	4	..	3	..	1	3	7	276	
November ..	56	47	23	86	1	7	1	17	5	15	6	1	3	..	1	1	..	2	5	277	
December ..	61	62	13	44	3	7	1	20	8	8	6	1	2	..	4	3	53	1	1	1	1	299	
Totals ..	829	802	195	904	16	148	24	233	110	142	62	17	37	4	23	21	72	44	44	7	24	1	* 3,715	
Totals, 1931 ..	1,304	1,327	161	1,109	22	25	247	233	160	133	84	21	59	9	33	12	24	31	31	1	13	2	1	1	1	..	* 5,013	
Totals, 1930 ..	2,244	1,440	149	1,244	30	12	104	322	157	162	93	30	52	7	36	15	18	68	68	5	6	3	1,040 7,238	

* Ceased to be a notifiable disease in January, 1931.