

*Summary of the Average Percentages from 9th January to 20th March,  
1913.*

Number for Factory Brand.	Moisture.	Fat.	Casein and other Solids.	Age, and Notes at Sampling.			
				Days old.	Body and Texture.		Remarks.
					Points.		
1 .. ..	35.60	34.60	29.80	15.6	28.1	..	
2 .. ..	36.03	34.67	29.30	21.5	28.3	..	
3 .. ..	35.52	34.33	30.45	22.3	28.8	..	
4 .. ..	34.12	35.13	30.75	19.5	28.8	..	
5 .. ..	37.08	32.55	30.37	16.0	28.0	..	
6 .. ..	35.78	33.73	30.49	22.2	28.5	..	
7 .. ..	35.92	33.73	30.35	23.5	28.7	..	
8 .. ..	35.54	35.08	29.38	25.6	28.2	..	
9 .. ..	34.85	35.47	29.68	26.0	28.8	..	
10 .. .	34.50	34.80	30.70	23.3	28.5	..	
Average ..	35.47	34.40	30.13	21.5	28.5	..	

*Summary of the Average Percentages in First Part of this List from  
17th October to 24th December, 1912.*

Number for Factory Brand.	Moisture.	Fat.	Casein and other Solids.	Age, and Notes at Sampling.			
				Days old.	Body and Texture.		Remarks.
					Points.		
1 .. ..	37.17	32.67	30.16	..	..	..	
2 .. ..	37.43	33.40	29.17	..	..	..	
3 .. ..	36.98	33.80	29.22	..	..	..	
4 .. ..	36.08	33.80	30.12	..	..	..	
5 .. ..	37.41	33.20	29.39	..	..	..	
6 .. ..	36.75	33.40	29.85	..	..	..	
7 .. ..	36.87	34.33	28.80	..	..	..	
8 .. ..	36.48	33.93	29.59	..	..	..	
9 .. ..	37.03	33.40	29.37	..	..	..	
10 .. ..	36.25	34.53	29.22	..	..	..	
Average ..	36.85	33.64	29.51	..	..	..	
General average	36.16	34.02	29.82			..	