

Table VII.—Summary of Carcass-quality Results.  
(Percentage Efficiency.)

	Series A.			Series B.			Series C.					
	Group 1.	Group 2.	Group 3.	Group 1.	Group 2.	Group 3.	Group 1.	Group 2.	Group 3.	Group 4.	Group 5.	
	Nil whole period ..	1 lb. whole period 140 lb.	2 lb. whole period 260 lb.	Nil whole period ..	¾ lb. whole period 130 lb.	1½ lb. whole period 200 lb.	Nil whole period ..	1 lb. whole period(1) 120 lb.	1 lb. early, nil late(2) 40 lb.	35 lb.—140 lb.	2 lb. early, 1 lb. late(1),(3) 160 lb.	Nil early, 1 lb. late(3) 85 lb.
Rate of meal-supplementing per 100 lb. live-weight per day	..	..	..	..	..	..	..	..	..	..	..	..
Meal used per 100 lb. dressed-weight gain	..	..	..	..	..	..	..	..	..	..	..	..
Weight range increase covered (dressed weight)	..	..	..	..	..	..	..	..	..	..	..	..
Number of pigs	10	10	10	12	12	12	8	8	8	8	8	8
	Percentage.	Percentage.	Percentage.	Percentage.	Percentage.	Percentage.	Percentage.	Percentage.	Percentage.	Percentage.	Percentage.	Percentage.
<i>Carcass Efficiency.</i>	94	95	93	87	94	87	51	45	54	44	55	55
* Streak	..	..	..	..	..	..	..	..	..	..	..	..
Eye of loin	..	..	..	..	..	..	..	..	..	..	..	..
Proportion of fat	..	..	..	..	..	..	..	..	..	..	..	..
Length of loin	..	..	..	..	..	..	..	..	..	..	..	..

(1) Whole period : 50 lb. — 200 lb. live-weight. (2) Early period : 50 lb. — 120 lb. live-weight. (3) Late period : 120 lb. — 200 lb. live-weight.  
\* Series A and B; Evaluation by measurement. Series C : Evaluation by standard photographs.