

# Weed Seeds In Agricultural Seed

(Sixth Series).


By

E. O. C. HYDE,  
Seed Analyst, Seed-Testing Station,  
Palmerston North.

The preceding five series of illustrated weed seeds appeared in the January, May, June, July, and August issues respectively of the "Journal."




*POLYDORA MONSPELIENSIS*: BEARD-GRASS. Straw coloured. Occurs rather infrequently in cocksfoot, crested dogtail, and ryegrass. An annual grass widely distributed, especially in coastal regions.




*FESTUCA ARUNDINACEA*: TALL FESCUE. Straw coloured. Occurs with medium frequency in cocksfoot and rarely in ryegrass. A coarse and rather unpalatable grass common in wet pasture land. The seed is very subject to ergot attack.




*PUCGINELLIA FASCICULATA*. Colour yellow to grey. Occurs frequently in creeping bent seed and rarely in *Lotus major* and white clover. A perennial grass often abundant on wet, salty land.



*ECHINOCHLOA CRUSGALLI*: BARNYARD-GRASS. Straw coloured. Occurs infrequently in cocksfoot, red clover, and mangold. An annual weed.



*HOLCUS MOLLIS*: CREEPING FOG OR SOFT FOG. Light straw colour. Occurs rarely in cocksfoot. A perennial grass spreading by underground stems.



*PHALARIS MINOR*: LESSER CANARY-GRASS. Colour grey. Occurs with medium frequency in white clover. An annual weed of little importance.