
Farmers Don't Feed Your Barns to the Borer!

Unprotected farm buildings are an invitation to wood-eating insects and rot producing fungi. The most successful way of preventing infestation or of controlling the pests is the use of pentachlorphenol as applied by BORACURE.




When BORACURE treatment is applied to income-earning buildings the cost is deductible in tax returns, as maintenance.

Protect your investment! Have a Boracure expert inspect your property thoroughly. There is no fee or obligation but quick action now could save you hundreds of pounds later. Get in touch with . . .

217 LAMBTON QUAY, WELLINGTON. 'Phone 40-050.

Branches Throughout the Country.





TUBERCULOUS MILKisa CAUSE OF DISEASE

In New Zealand in 1945, 187 people died from non-pulmonary tuberculosis — that is, tuberculosis of the brain, the bowels, the glands, the bones and the joints. Many of these deaths could have been prevented, because many were due to milk-borne tuberculosis.

Fatal infection can take place in childhood through drinking impure milk.

Milk-borne tuberculosis can be prevented -

- 1. By boiling the milk (which kills the tubercle bacilli and alters the composition of the milk)
- 2. By pasteurising all milk (which also kills the tubercle bacilli but does not impair the nutritive value).

If you cannot get pasteurised milk, here is an easy home method of pasteurising it yourself; fill the lower part of a double boiler with water, place the milk in the upper part, cover, and heat until the water boils; allow it to boil for 3 minutes; remove and cool immediately by placing the upper part (which contains the milk) in a large vessel of cold water; change the water every few minutes until the milk is as cold as the water.

TUBERCULOSIS CAN SPREAD FROM COWS
TO HUMANS, BUT NOT IF YOU



ISSUED BY THE DEPARTMENT OF HEALTH

Keep this announcement for future reference.