

Lyreidus fessor Bennett, 1964: 24–6, figs. 5–9, 106 (type locality: North of Whale Island, Bay of Plenty, New Zealand, 67fms; holotype (male) in Canterbury Museum, Christchurch).

TYPES: Dr L. B. Holthuis has kindly informed me that the material on which De Haan based his original description of *L. tridentatus* comprises 6 syntypes. On his advice I select the most complete of these as LECTOTYPE: A female, c.l. 40mm, Japan, 1823–1835, P. F. von Siebold and H. Burger, Crust. reg. no. D.23017, a dry specimen with the mouthparts dissected out and pasted on cardboard.

PARALECTOTYPES: Two dry specimens with same data as lectotype; 3 dry specimens, Japan, 1830–1835, H. Burger.

LOCALITIES PREVIOUSLY REPORTED

Japan: "Japan" (De Haan). Kada Bay (Miers). S. of Inuboe-zaki, 196 m. ("Misago" Sta. 113); E. of Bosyu, 117–269 m. ("Misago" Sta's 5, 103, 104, 105) (Yokoya). Tokyo Bay (Doflein, 1902; Parisi; Balss). Sagami Bay (Ortmann; Parisi). Misaki, Sagami Bay, 100 m., 200–300 m. (Parisi; Balss; Sakai, 1937). Dzushi, 130 m., and Ito, Sagami Bay (Balss). Manazuru and Hayama, Sagami Bay (Sakai 1937). Enoshima, Sagami Bay (Parisi). Off Enoshima, 30–80 m., off Kameki reef and off Mito-hama, 30–50 m., Sagami Bay (Sakai, 1965). Suruga Bay, 123 m. ("Misago" Sta. 286); S. of Lake Hamana, 172 m. ("Misago" Sta. 195); S. of Sima, Mie-ken, 207 m. ("Misago" Sta. 199); S. of Hamazima, Mie-ken, 93 m. ("Misago" Sta. 365); N.E. of Owrase, Mie-ken, 183 m. ("Misago" Sta. 201); W. of Siwo-misaki, 165 m. ("Misago" Sta. 210) (Yokoya). Tosa Bay (Sakai, 1937). N.E. of Asizurizaki, 223 m. ("Misago" Sta. 226); S. of Kōti, 97 m. and 126 m. ("Misago" Sta's 341 and 406); Bungo Strait, 241 m. ("Misago" Sta. 304) (Yokoya). Nagasaki (Sakai, 1934, 1937). S. of Goto Is., 238 m. ("Misago" Sta. 238); Toyama Bay, 135 m. ("Misago" Sta. 580) (Yokoya). Fukuura (Balss). Off Tori-shima (Ryukyu Islands), 100 fms (Miyake).

South China Sea: Hong Kong (Wood-Mason, 1887).

South Pacific Ocean: Off Kandavu, Fiji Is., 210 fms ("Challenger" Sta. 174) (Henderson). New Caledonia (Ortmann).

Central Pacific Ocean: Hawaii (Tinker).

Australia: Pt Stephens (Haswell). Off Newcastle, 45–50 fms (Ward). Newcastle Bight, 5–7 miles off shore, 26–40 fms ("Thetis" Sta. 22); 64–8 miles off Newcastle, 42–48 fms ("Thetis" Sta. 25) (Whitelegge). Outside the heads of Pt Jackson (Haswell). Off Pt Jackson, 30–35 fms (Henderson). Off Wata Mooli, 3½–4 miles offshore, 54–59 fms ("Thetis" Sta. 57) (Whitelegge).

New Zealand: Outside Great Barrier I., 120 fms (Chilton). Near the entrance to Hauraki Gulf, probably in about 20–25 fms (Powell, 1947). W. of Aldermen Is., Bay of Plenty, 50 fms; between Mayor I. and Motiti I., Bay of Plenty, 70–100 fms; N.E. of Motiti I., Bay of Plenty, 49 fms; 7 miles S.E. of Whale I., Bay of Plenty (Powell, 1949). Off White I., Bay of Plenty, 113–120 fms; 14 miles E. of Whale I., Bay of Plenty, 105 fms (Dell, 1963). N. of Whale I., Bay of Plenty, 67 fms (Bennett). Three miles N.E. of Hicks Bay, 100–120 fms (Dell, 1963a). Waikawa Point, near C. Runaway, 60 fms (Powell, 1949). "Ex Napier trawlers" (Dell, 1956). "Wellington area, from dogfish stomach" (Richardson and Krefft).

MATERIAL EXAMINED: A total of 243 specimens, as follows—Japan: "Japan", Reiter, 1894, 1 ♂, ca 31mm (SM unreg.—photo only). Japanese Seas, pres. T. Sakai before April 1934, 1 ♂, 39.9mm (AM P.10499), 1 ♀, 41.7mm (AM P.10500). Off Honshu, 65 fms, 7/5/1900, "Albatross" Sta. 3740, 1 ♂, 17.0mm (USNM 57686). Misaki, don. A. S. Pearse, 1 ♂, 28.3mm (USNM 63689). Sagami Bay, Doderlein, 1882, 2 ♀♀, ca 29, 44mm (SM unreg.—photos only).

South China Sea: S. of Hong Kong, 21°04.5' to 21°05'N., 114°30.5'E., 46 fms, mud and coarse sand, Agassiz trawl, 25/7/1965, FHK Sta. 18, 1 juv., 10.9mm (AM P.15784). S.E. of Hong Kong, 21°14'N., 115°18'E., 62 fms, Agassiz trawl, 13/8/1965, FHK Sta. 21, 1 ♀, 28.9mm (AM P.15785). S.E. of Hong Kong, 21°15.2'N., 115°16.4'E., 63 fms, Agassiz trawl, 13/8/1965, FHK Sta. 19, 1 ♀, 24.6mm (AM P.15786). N.E. of Bunguran, 5°41'N., 110°07' to 110°10'E., 103–131 fms, muddy sand, Granton trawl, 6/11/1964, FHK Sta. 41, 1 empty carapace with sternum and mandibles, 22.2mm (AM P.15783).

Central Pacific Ocean: Hawaii, T. Richert, 1 ♀, 20.3mm (WAM 351–67). Hawaii, 200 fms, T. Richert, 2 ♂♂, 2 ♀♀, 16.7–41.7mm (WAM 352–67). Off Sand I., Oahu, Hawaii, 60 fms, 18/7/1959, "Pele" Expedition Sta. 168, 1 juv., 1 ♀, 13.2, 27.8mm (BPBM 6796). Off Waikiki, Oahu, Hawaii, 90–102 fms, "Pele" Expedition, 8/4/1959, 2 juvs., 3.5, 13.5mm (BPBM 6797).

South Pacific Ocean: New Caledonia, Krieger collection, 1864, 2 ♂♂, 37.7, 39.6mm (SM unreg.).