

**Paralia** Heiberg

**P. sulcata** (Ehrenb.) Cleve (*Melosira sulcata* (Ehrenb.) Kütz.)—Widely distributed. 14 records.

**Planktoniella** Schütt

\***P. florea** Wood—Waitarere Beach, CC.

\***P. sol** (Wall.) Schütt—Hauraki Gulf, C1; Hawke Bay, C2; Castle Point, C1; W of Hawera, C2; New Plymouth/Kawhia, C2.

**Podosira** Ehrenb.

**P. hormoides** (Mont.) Kütz. (*Hyalodiscus hormoides* Kütz.)—Lyall Bay, P.

\***P. stelliger** (Bail.) Mann (*Hyalodiscus stelliger* Bail.; *H. maculata* W. Sm.)—Goat Island Bay, Leigh, T; Hauraki Gulf, C1; Auckland Harbour, CW1; Bay of Plenty, C2; Lyall Bay, P (as both *H. maculatus* and *H. stelliger*); W of Hawera, C2; Lyttelton Harbour, C2; Foveaux Strait, P.

**Pyxidicula** Ehrenb.

\***P. cruciata** Ehrenb.—Auckland Harbour, CW1; Whangateau Harbour, T.

**Skeletonema** Grev.

\***S. costatum** (Grev.) Cleve—Widely distributed. 15 records.

**Stephanopyxis** Ehrenb.

\***S. orbicularis** Wood—Widely distributed. 16 records. (TYPE) The type locality given in WCC is “planktonic in New Zealand coastal waters”.

\***S. turris** (Grev. & Arnott) Ralfs—Widely distributed. 14 records.

**Thalassiosira** Cleve<sup>7</sup>

\***T. condensata** Cleve—Widely distributed. 11 records.

\***T. decipiens** (Grun.) Jørg.—Widely distributed. 12 records.

\***T. hyalina** (Grun.) Gran—Widely distributed. 16 records.

\***T. polychorda** (Gran) Proskina-Lavrenko (*Coscinosira polychorda* (Gran) Gran)—Hauraki Gulf, C1.

\***T. rotula** Meunier—Widely distributed. 8 records.

\***T. subtilis** (Ostenf.) Gran—NE coast, CW1.

## Family HEMIDISCACEAE

**Hemidiscus** Wall.

\***H. cuneiformis** Wall.—N of New Zealand, CW1; Hawke Bay, C2.

## Family ACTINODISCACEAE

**Actinoptychus** Ehrenb.

\***A. adriaticus** Grun.—Waitarere Beach, CC; New Plymouth/Kawhia, C2.

**A. punctatus** (Ehrenb.) Castr. (non *A. punctatus* Bail.)—Lyall Bay, P.

\***A. senarius** (Ehrenb.) Ehrenb. (*Actinocyclus undulatus* Kütz.)—Widely distributed. 12 records.

\***A. splendens** (Shad.) Ralfs ex Pritch.—Goat Island Bay, Leigh, T; Lyall Bay, P; Otago Peninsula, C2.

**A. vulgaris** Schum.—Smith's Bay, Dunedin, W2; Bruce's Rocks, Dunedin, W2; Goat Island, Dunedin, W2.

**A. sp.**—Bruce's Rocks, Dunedin, W2.

**Asteromphalus** Ehrenb.

\***A. flabellatus** (de Bréb.) Grev.—Hauraki Gulf, WCC, C1, C2; Bay of Plenty, C2; Hawke Bay, C2; Lyall Bay, P.

\***A. hookeri** Ehrenb.—Waitarere Beach, CC; Cook Strait, C1.