

APPENDIX 2: FOSSIL SEEDS FROM THE LINDALE SECTION,
PARAPARAUMU

By C. R. LENNIE

New Zealand Geological Survey*

The samples from N157/520 examined for fruits and seeds weighed from 10 to 21lb. They yielded poor seed floras. Ten unidentified seed types were recorded besides those in Table II. The stratigraphic position of the samples, registered as B839-45, are shown in Fig. 3.

TABLE II.

	<i>Montia fontana</i> L.	<i>Hydrocotyle</i> sp.	<i>Rubus</i> sp.	<i>Oxalis</i> sp.	Compositae sp. 1	Compositae sp. 2	<i>Haloragis ? uniflora</i> Kirk	<i>Potamogeton</i> sp.	<i>Juncus</i> sp.
B839 T	x	x	-	-	x	-	-	-	x
B840 U	-	-	x	-	-	-	-	x	x
B841 V	x	x	-	-	-	-	x	x	x
B842 W	x	x	-	x	-	-	-	x	x
B843 X	x	x	-	-	-	x	-	x	x
B844 Y	x	x	-	x	-	-	-	x	x
B845 Z	-	-	-	-	-	x	-	x	x

The samples were also examined for insect remains; a number of insect fragments were found, but none has yet been identified.

* Present address: Shell Development (Australia) Pty. Ltd., Melbourne, Victoria, Australia, 3001.