

TABLE II.—Dimensions of *Eutrephoceras* sp.

	Diameter	Width (W)	Height of Whorl (H)	W / H	Distance from venter (V)	Median height of Whorl (M)	M / V
GS 13	61.5	52.0	45.0	1.15	7.2	13.6	1.88
C 697	165.0	128.0*	121.0*	1.07	47.5	60.5	1.28

\* Calculated.

REMARKS: Both specimens have a similar shape, but C 697 has a wider umbilicus and the siphuncle is closer to the dorsum than in GS 13.

## ACKNOWLEDGMENTS

The writer is indebted to Dr A. G. Beu and Dr C. A. Fleming, New Zealand Geological Survey, for constructive criticism of the manuscript.

## REFERENCES

- BEU, A. G., 1968. A specimen of the Nautiloid *Aturia* from the Kapitean Stage (Uppermost Miocene) of New Zealand. *N.Z. Jl. Geol. Geophys.* 11(1): 161-5.
- BLANFORD, H. F., 1861. In BLANFORD, H. F.; STOLICKA, F. The fossil Cephalopoda of the Cretaceous rocks of Southern India; *Belemnitidae-Nautilidae Mem. geol. Surv. India (Palaeont. Indica)*, Ser. 1: 1-40.
- FLEMING, C. A., 1945. Some New Zealand Tertiary cephalopods. *Trans. R. Soc. N.Z.* 74: 411-18.
- GLENISTER, B. F.; MILLER, A. K.; FURNISH, W. M., 1956. Upper Cretaceous and early Tertiary nautiloids from Western Australia. *J. Paleont.* 30: 492-503.
- HAAS, O.; MILLER, A. K., 1952. Eocene nautiloids of British Somaliland. *Amer. Mus. nat. Hist. Bull.* 99: 317-54.
- JOHNSTON, M. R. (in preparation). Sheet N 159—Tinui. "Geological Map of New Zealand 1 : 63,360." New Zealand Department of Scientific and Industrial Research.
- KUMMEL, B., 1956. Post-Triassic nautiloid genera. *Bull. Mus. comp. Zool. Harv.* 114(7): 324-494.
- 1964. Nautiloidea—Nautilida. *Treatise on Invertebrate Paleontology, Pt K Cephalopoda, Nautiloidea.*
- MCGOWRAN, B., 1959. Tertiary nautiloids (*Eutrephoceras* and *Cinomia*) from South Australia. *J. Paleont.* 33: 435-48.
- MCKELLAR, I. C., 1969. Sheet S 169—Winton. "Geological Map of New Zealand 1 : 63,360." New Zealand Department of Scientific and Industrial Research.
- MATSUMOTO, T., 1967. A Cretaceous nautiloid from Urakawa, Hokkaido. *Jap. J. Geol. Geog. Trans.* 38: 163-9.
- MILLER, A. K., 1947. Tertiary nautiloids of the Americas. *Mem. geol. Soc. Am.* 23: 1-234.
- SPENGLER, E., 1910. Untersuchungen über die sudindische Kreideformation: Die Nautiliden und Belemniten des Trichinopolydistrikts. *Beitr. Palaont. Osterreich-Ungarns und Oriens* 23: 125-57.

M. R. JOHNSTON,  
N.Z. Geological Survey,  
Nelson,  
New Zealand.