

genera of which nine are indigenous; and the Eriococcidae (Hoy, 1962) in which the 75 known species are distributed among nine genera of which seven are indigenous.

## LITERATURE CITED

- BERLESE, A., 1882-1893. Acari, Myriapoda et Scorpiones hucusque in Italia Reperta. Ordo Prostigmatæ (Trombidiidae). Patavii, Portici.
- 1910. Acari nuovi. Maniplus V. *Redia* 6: 199-234.
- CANESTRINI, G., 1889. Prospetto del 'Acarofauna Italiana. Famiglia dei Tetranychini. *Atti Ist. veneto Sci.* 7(6): 491-537.
- CHAUDRHI, W. M., 1965. New mites of the genus *Ledermuelleria*. *Acarologia* 7: 467-486.
- COLLYER, E., 1964. Phytophagous mites and their predators in New Zealand orchards. *N.Z. J. agric. Res.* 7: 551-568.
- COTTIER, W., 1934. The natural enemies of the European Red Mite in New Zealand. *N.Z. J. Sci. Tech.* 16: 68-80.
- EHARA, S., 1962. Notes on some predatory mites (Phytoseiidae and Stigmaeidae). *Jap. J. appl. Ent. Zool.* 6: 53-60.
- GIVEN, B. B., 1952. A revision of the Melolonthinae of New Zealand. Part I: the adult beetles. *N.Z. D.S.I.R. Bull.* 102: 1-137.
- GONZALEZ, R. H., 1963. Four new mites of the genus *Agistemus* Summers. *Acarologia* 5: 342-350.
- 1965. A taxonomic study of the genera *Mediolata*, *Zetzellia* and *Agistemus* (Acarina: Stigmaeidae). *Univ. Calif. Publ. Ent.* 41: 64 pp.
- GRANDJEAN, F., 1944. Observations sur les acariens de la famille des Stigmaeidae. *Archs. Sci. phys. nat.* 26: 103-131.
- HABEEB, H., 1958. New mites from New Brunswick. *Leaflet. Acadian Biol.* 18: 1-4.
- 1961. Walter Vincent Powers, Nobel Fellow. *Ibid.* 22: 1-6.
- HALBERT, J. N., 1920. The Acarina of the seashore. *Proc. roy. Irish Acad.* 35: 106-152.
- HIRST, S., 1926. Report on the acari found on or associated with sandflies in India. *Indian J. med. Res.* 13: 1023-1026.
- HOY, J. M., 1962. Eriococcidae (Homoptera: Coccoidea) of New Zealand. *N.Z. D.S.I.R. Bull.* 146: 219 pp.
- KOCH, C. L., 1835-1844. Deutschlands Crustaceen, Myriapoden und Arachniden. Heft 1-40. Regensburg.
- LAMB, K. P., 1952. A preliminary list of New Zealand Acarina. *Trans. roy. Soc. N.Z.* 79: 370-375.
- LEE, K. E., 1959. The earthworm fauna of New Zealand. *N.Z. D.S.I.R. Bull.* 130: 486 pp.
- MITRA, R. D., and MITRA, S. D., 1953. A new species of *Raphignathus* associated with *Phlebotomus* in India. *Z. Parasit. Kde.* 15: 429-432.
- OUDEMANS, A. C., 1923a. Acarologische Aanteekeningen LXX. *Ent. Ber., Amst.* 6(129): 138-144.
- 1923b. Acarologische Aanteekenigen LXXI. *Ent. Ber., Amst.* 6(130): 145-155.
- 1927. Acarologische Aanteekenigen LXXXVIII. *Ibid.* 7(158): 263.
- SUMMERS, F. M., 1960a. *Eupalopsis* and eupalopsellid mites (Acarina: Stigmaeidae, Eupalopsellidae). *Fla. Ent.* 43: 119-138.
- 1960b. Several stigmaeid mites formerly included in *Mediolata* redescribed in *Zetzellia* Ouds., and *Agistemus*, new genus. *Proc. ent. Soc. Wash.* 62: 233-247.
- 1962. The genus *Stigmaeus* (Acarina: Stigmaeidae). *Hilgardia* 33: 491-537.
- 1964. Three uncommon genera of the mite family Stigmaeidae (Acarina). *Proc. ent. Soc. Wash.* 66: 184-192.