

*Cingula (Hyala)* H. and A. Adams, 1852 (*H. vitrea* (= *Turbo vitreus*) Montagu, 1803; monotypy).

*Cingula (Setia)* H. and A. Adams, 1852 (*Rissoa pulcherrima* Jeffreys, 1848; s.d. Kobelt, 1878).

*Elachisina* Dall, 1918 (*E. grippi* Dall, 1918; monotypy).

*Falsicingula* Habe, 1958 (*Cingula kuriensis* Pilsbry, 1905; o.d.).

Most of the North American species classified as *Cingula* should probably be located under *Falsicingula*.

*Mistostigma* Berry, 1947 (*M. punctulum* Berry, 1947; o.d.).

? *Nanadoma* Laseron, 1956 (*N. imitoris* Laseron, 1956; o.d.).

This genus probably does not belong in the Rissoidae.

*Onoba* H. and A. Adams, 1852, nom. nov. pro *Turbanilla* Leach, 1847, non Risso, 1826 (*O. striata* (= *Turbo striatus* J. Adams, 1797, non Da Costa, 1778 = *Turbo semicostatus* Montagu, 1803); monotypy).

The shell of *Onoba* has a different "style" and aperture from *Cingula* and generic distinction seems justified.

*Onoba (Onoba)* s.s.

*Onoba (Microdochus)* Rehder, 1943 (*M. floridanus* Rehder, 1943; o.d.).

*Ovirissoa* Hedley, 1916 (*Rissoa adarensis* Smith, 1902; o.d.).

*Ovirissoa (Ovirissoa)* s.s.

*Ovirissoa (Paranoba)* Laseron, 1950 (*P. subquadrata* Laseron, 1950; o.d.).

Placed here tentatively, *Austrorissopsis* Ponder is possibly related.

*Parashiela* Laseron (*P. ambulata* Laseron, 1956; o.d.).

*Parvisetia* Monterosato, 1884 (*Rissoa scillae* Aradas and Benoit, 1870, ex Seguenza MS; monotypy).

*Parvisetia* appears to be a good genus and quite distinct from *Setia* with which it was placed as a synonym by Coan (1964).

Synonym: *Obtusella* Cossmann, 1921 nom. nov. pro *Cingulina* Monterosato, 1884, non A. Adams, 1860 (*Rissoa obtusa* Centraine, 1842 (non Brown, 1841) ? = *R. soluta* Philippi, 1844; monotypy).

*Plagystila* Fischer in De Folin and Périer, 1872, pro *Plagiostyla* Fischer in De Folin and Périer, 1871, nom. nud. (*P. asturiana* Fischer in De Folin and Périer, 1872; monotypy).

*Powellisetia* Ponder, 1965 (*Rissoa porcellana* Suter, 1908; o.d.).

*Pseudosetia* Monterosato, 1884 (*Rissoa turgida* Jeffreys, 1870; s.d. Crosse, 1885).

The operculum of this genus is thin and horny and, therefore, it is not related to *Barleeia* as Coan has indicated.

*Putilla* A. Adams 1867 (*P. lucida* A. Adams, 1867; monotypy).

*Rissopsis* Garrett, 1873 (*R. typica* Garrett, 1873; monotypy).

The following subgenera are only tentative.

*Rissopsis (Rissopsis)* s.s.

*Rissopsis (Antinodulus)* Cossman in Cossman and Peyrot, 1919 (*Bulimus globulus* Grataloup, 1828; o.d.).

*Rissopsis (Peringiella)* Monterosato, 1878 (*Rissoa laevis* Monterosato, 1877; monotypy) (see Ponder, 1965c).

*Rissopsis (Pisinna)* Monterosato, 1878 (*Rissoa punctulum* Philippi, 1836 = *Helix glabrata* Mühlfeld, 1824; s.d. Cossmann, 1921).