



FIG. 1.—*Megacolabus sculpturatus* Broun. a, left fore tibia; b, right hind tibia; c, left antenna; d, lateral view; e, dorsal view.

KEY TO THE SPECIES OF MEGACOLABUS

- | | | | | | |
|---|-------|-------|-------|-------|------------------------------|
| 1. Funicle of antenna with joint 1 equal to joint 2 | | | | | 2 |
| Funicle of antenna with joint 1 longer than joint 2 | | | | | 5 |
| Funicle of antenna with joint 1 shorter than joint 2 | | | | | <i>M. harrisi</i> (Brookes) |
| 2. Pronotum more or less quadrate; sides gently rounded or parallel | | | | | 3 |
| Pronotum distinctly transverse, ratio 0.77; sides strongly rounded | | | | | <i>M. sculpturatus</i> Broun |
| 3. Elytral punctures indistinctly seriate; elevations conspicuous; larger species | | | | | 4 |
| Elytral punctures in distinct series; elevations feeble; smaller species | | | | | <i>M. pteridosus</i> n.sp. |
| 4. Elytral intervals 4 and 6 with uneven sculpture and clothing; antennae with scape slender, abruptly clavate at apex | | | | | <i>M. decipiens</i> Marshall |
| Elytral intervals 4 and 6, in apical half, smooth and evenly clothed; antennae with scape robust, expanding gradually to apex | | | | | <i>M. garviensis</i> n.sp. |
| 5. Elytra much wider than pronotum, ratio 1.62; strongly sculptured species | | | | | <i>M. obesus</i> n.sp. |
| Elytra slightly wider than pronotum, ratio 1.33; sculpture relatively feeble | | | | | <i>M. reductus</i> Marshall |